

Home Safety Fact Sheet (2014)

Fatalities

Suffocation: 1,169 children ages 19 and under died from suffocation in 2011.¹

- 77% of children (896) who suffocated were under the age of 1 year.¹
- 59% of children (692) who suffocated were boys.¹
- In 2010, 88 children died from accidental hanging and strangulation. 147 children died from choking on food or another object.²

Falls: 138 children ages 19 and under died in falls in 2011.¹


- 54% of children (75) who died in falls were ages 15 to 19.¹
- 78% of children (108) who died in falls were boys.¹

TV tip-overs: On average, 17 children die from TV tip-overs each year.³

Toys: 11 toy-related deaths in children under the age of 15 years were reported to the U.S. Consumer Product Safety Commission in 2012.⁴

Button batteries: 3 children under the age of 6 died from button battery ingestion and were reported to poison control centers in 2013.⁵

1999-2011 Fall and Suffocation Fatalities and Death Rates Among Children Ages 19 and Under



Injuries

Suffocation: 24,860 children were seen in emergency departments for nonfatal suffocation or inhalation in 2012.¹

Falls: 2,812,190 children were seen in emergency departments for nonfatal falls in 2012.¹

TV tip-overs: 12,800 kids aged 19 and under were seen in emergency departments for injuries caused by televisions tipping over in 2011.⁶

Toys: 192,000 children under the age of 15 years were seen in emergency departments for injuries caused by toys in 2012.⁴

Button batteries: Poison control centers reported 2,759 cases of children ages 19 and under swallowing button batteries in 2013.⁵

Additional Statistics

Suffocation

- 70% of suffocation deaths among infants were from accidental suffocation or strangulation in bed in 2010.²
- In 2010, cribs and playpens were responsible for more than 20 percent of all nursery product-related emergency department-treated injuries among children ages 5 and under.⁷
- The majority of childhood suffocation, choking and strangulation incidents occur in the home.^{8,9}

Falls

- Window falls account for approximately 8 deaths and 3,300 injuries among children ages 5 and under annually.¹⁰
- The risk of a child being injured as a result of a fall at home is twice the risk as at child-care.¹¹
- Window falls occur more frequently in large urban areas and low-income neighborhoods.^{12,13}
- In New York City and Boston, education and window guard distribution programs resulted in a 96 percent reduction in the incidence of window falls over 10 years.¹⁴

TV/Furniture tip-overs

- The estimated number of emergency room visits for kids aged 19 and under has increased 31% over the last decade, from 9,800 in 2002 to 12,800 in 2011.⁶
- Because of its weight, a 36-inch CRT television falling three feet creates the same momentum as a 1-year-old child falling 10 stories.¹⁵
- Approximately \$8.34 million is spent in medical costs each year to treat children aged 19 and under in emergency departments for injuries from TV tip-overs.^{6,16}
- 7 out of 10 (72%) of children who are injured by a TV tipping over are 5 years old or younger.⁶

Toys

- 45% of toy-related injuries are to the head and face.⁴
- One-third of toy-related injuries occur in children under the age of 5 years.⁴
- The category of toy responsible for the most injuries in children under the age of 15 years are nonmotorized scooters, accounting for 27% of injuries.⁴

Button batteries



- In two hours, a button battery ingestion can cause severe, often irreparable damage to a child. Most severe complications effect the esophagus.¹⁷
- The average age of a child seen in the emergency department for swallowing a button battery is 3.9 years.¹⁸
- Button batteries are involved in 84% of all battery-related emergency department visits, for which a battery type was identified.¹⁹
- When a button battery gets stuck in a child’s throat, the saliva triggers an electrical current that causes a chemical reaction which burns the esophagus. Damage can continue even after the battery is removed.¹⁷
- The number of serious injuries or deaths as a result of button batteries has increased more than 9-fold in the past decade.⁵

References

- ¹ Centers for Disease Control and Prevention, National Center for Injury Prevention and Control. Web-based Injury Statistics Query and Reporting System (WISQARS). National Center for Injury Prevention and Control Website. Unintentional falls and suffocation fatalities and injuries, children ages 19 and under. Available from: <http://www.cdc.gov/injury/wisqars/index.html>. Fatalities accessed June 2014, nonfatal injuries accessed March 2013.
- ² Centers for Disease Control and Prevention. WONDER Query System. Available at: <http://wonder.cdc.gov/>. Accessed March 27, 2013 (fatal) and October 29, 2013 (nonfatal).
- ³ Gipson K. Product Instability or Tip-Over Injuries and Fatalities Associated with Televisions, Furniture, and Appliances: 2012 Report. U.S. Consumer Product Safety Commission, November 2012.
- ⁴ Tu Y. Toy-Related Deaths and Injuries Calendar Year 2012. U.S. Consumer Product Safety Commission, November 2013.
- ⁵ National Capital Poison Center. Button Battery Ingestion Statistics. Available at: <http://www.poisson.org/battery/stats.asp>. Accessed June 18, 2014.
- ⁶ Ferguson RW, Mickalide AD. A Report to the Nation on Home Safety: The Dangers of TV Tip-Overs. Washington, DC: Safe Kids Worldwide, December 2012.
- ⁷ Chowdhury RT. Nursery product-related injuries and deaths among children under age five. Bethesda, MD: Consumer Product Safety Commission, Division of Hazard Analysis; December 2011.
- ⁸ Reilly JS, Walter MA. Consumer product aspiration and ingestion in children: analysis of emergency room reports to the National Electronic Injury Surveillance System. *Ann Otol Rhinol Laryngol*. 1992. 101: 739-41.
- ⁹ Children’s Hospital Boston. Airway obstruction. Children’s Hospital Boston Website. Available at: <http://www.childrenshospital.org/az/Site772/mainpageS772P1.html>. Accessed October 13, 2011.
- ¹⁰ United States Consumer Product Safety Commission. CPSC: Parents, caregivers should consider safety before opening windows, April 8, 2011. Washington, DC: Office of Information and Public Affairs.
- ¹¹ Khambalia A, Joshi P, Brussoni M, Raina P, Morrongiello B, Macarthur C. Risk factors for unintentional injuries due to falls in children ages 0-6 years: a systematic review. *Inj Prev*. 2006. 12(6): 378-385.
- ¹² Stone KE, Lanphear BP, et al. Childhood injuries and deaths due to falls from windows. *J Urban Health: Bulletin of the NY Academy of Med*. 2000. 77(1): 26-33.
- ¹³ Committee on Injury and Poison Prevention. Falls from heights: windows, roofs and balconies. *Pediatrics*. 2001. 107(5): 1188-1191.
- ¹⁴ Harris VA, Rochette LM, Smith GA. Pediatric injuries attributable to falls from windows in the United States in 1990-2008. *Pediatrics*. 2011. 128(3): 455-462.
- ¹⁵ Bernard PA, Johnston C, Curtis SE, King WD. Toppled television sets cause significant pediatric morbidity and mortality. *Pediatrics*. 1998. 102(3):E32.
- ¹⁶ Centers for Disease Control and Prevention. WISQARS Cost of Injury Reports, 2005. Available at <http://www.cdc.gov/injury/wisqars/cost/cost-learn-more.html>. Accessed 14 November 2012.
- ¹⁷ National Capital Poison Center. Mechanism of Injury. Available at: <http://www.poisson.org/battery/mechanism.asp>. Accessed March 27, 2013.
- ¹⁸ Sharpe SJ, Rochette LM, Smith GA. Pediatric battery-related emergency department visits in the United States, 1990-2009. *Pediatrics*. 2012. 129(6): 1111-7.
- ¹⁹ CDC MMWR. Injuries from batteries among children aged <13 years—United States, 1995-2010. *MMWR*. 2012. 61(34):661-666.

Last updated June 2014. If you have a question about this factsheet, please call 202-662-0600.

